
Artificial urinary sphincter in men: procedure-specific information

What is the evidence base for this information?

This leaflet includes advice from consensus panels, the British Association of Urological Surgeons, the Department of Health and evidence-based sources; it is, therefore, a reflection of best practice in the UK. It is intended to supplement any advice you may already have been given by your GP or other healthcare professionals. Alternative treatments are outlined below and can be discussed in more detail with your Urologist or Specialist Nurse.

What does the procedure involve?

The artificial urinary sphincter consists of three components. One part is a circular cuff that is placed around the waterpipe (urethra). This cuff is connected to a small pump that sits in the scrotum and also connected to small fluid-filled balloon that sits in the abdominal wall.



What are the alternatives to this procedure?

Incontinence into a pad, a urethral catheter or a male sling (the last option must be performed in the context of a clinical trial).

What should I expect before the procedure?

You will usually be admitted on the same day as your surgery. You will normally receive an appointment for pre-assessment, approximately 14 days before your admission, to assess your general fitness, to screen for the carriage of MRSA and to perform some baseline investigations. After admission, you will be seen by members of the medical team which may include the Consultant, Specialist Registrar, House Officer and your named nurse.

You will be asked not to eat or drink for 6 hours before surgery and, immediately before the operation, you may be given a pre-medication by the anaesthetist which will make you dry-mouthed and pleasantly sleepy.

Please be sure to inform your Urologist in advance of your surgery if you have any of the following:

- an artificial heart valve
- a coronary artery stent
- a heart pacemaker or defibrillator
- an artificial joint
- an artificial blood vessel graft
- a neurosurgical shunt
- any other implanted foreign body
- a prescription for Warfarin, Aspirin or Clopidogrel (Plavix®)
- a previous or current MRSA infection
- high risk of variant CJD (if you have received a corneal transplant, a neurosurgical dural transplant or previous injections of human-derived growth hormone)

What happens during the procedure?

Normally, a full general anaesthetic will be used and you will be asleep throughout the procedure. In some patients, the anaesthetist may also use an epidural anaesthetic which improves or minimises pain post-operatively. You will have a small incision in the area between the scrotum and anus (the perineum) and the cuff will be placed around the waterpipe. A separate incision will be made in the groin. Through this second wound, the pump will be placed in the scrotum and the balloon placed in the abdominal wall.

What happens immediately after the procedure?

You will be given fluids to drink from an early stage after the operation and you will be encouraged to mobilise as soon as you are comfortable to prevent blood clots forming in your legs. You will be given intravenous antibiotics through your vein (intravenous). You will usually be discharged home three days after your surgery.

Are there any side-effects?

Most procedures have a potential for side-effects. You should be reassured that, although all these complications are well-recognised, the majority of patients do not suffer any problems after a urological procedure.

Please use the check boxes to tick off individual items when you are happy that they have been discussed to your satisfaction:

Common (greater than 1 in 10)

- Blood in the urine , and temporary stinging when you urinate after the procedure
- In the long-term, mechanical failure of the device
- In the long-term, infection of the device requiring removal

Occasional (between 1 in 10 and 1 in 50)

- Urine infection
- Wound infection
- Temporary insertion of a bladder catheter
- Later failure of the device as the cuff becomes loose around the waterpipe (urethral atrophy)

Rare (less than 1 in 50)

- Erosion of the waterpipe through the waterpipe (urethra)

Hospital-acquired infection (overall risk for Addenbrooke's)

- Colonisation with MRSA (0.02%, 1 in 5,000)
- Clostridium difficile bowel infection (0.04%; 1 in 2,500)
- MRSA bloodstream infection (0.01%; 1 in 10,000)

(These rates may be greater in high-risk patients e.g. with long-term drainage tubes, after removal of the bladder for cancer, after previous infections, after prolonged hospitalisation or after multiple admissions)

What should I expect when I get home?

The device will be deactivated when you are discharged. You will be reviewed in clinic or on the ward at six weeks post-discharge when the device will be activated (by pressing a button on the pump within the scrotum).

What else should I look out for?

Are there any other important points?

You will be reviewed in outpatients to see how you have got on. It is likely you will be asked to complete a questionnaire on your symptoms.

Driving after surgery

It is your responsibility to ensure that you are fit to drive following your surgery. You do not normally need to notify the DVLA unless you have a medical condition that will last for longer than 3 months after your surgery and may affect your ability to drive. You should, however, check with your insurance company before returning to driving. Your doctors will be happy to provide you with advice on request.

Is there any research being carried out in this field at Addenbrooke's Hospital?

There is no specific research in this area at the moment but all operative procedures performed in the department are subject to rigorous audit at a monthly Audit & Clinical Governance meeting.

Who can I contact for more help or information?

Oncology Nurses

- Uro-Oncology Nurse Specialist
01223 586748
- Bladder cancer Nurse Practitioner (haematuria, chemotherapy & BCG)
01223 274608
- Prostate cancer Nurse Practitioner
01223 247608 or 216897 or bleep 154-548
- Surgical Care Practitioner
01223 348590 or 256157 or bleep 154-134

Non-Oncology Nurses

- Urology Nurse Practitioner (incontinence, urodynamics, catheter patients)
01223 274608 or 586748
- Urology Nurse Practitioner (stoma care)
01223 349800

Patient Advice & Liaison Centre (PALS)

- Telephone
+44 (0)1223 216756 or 257257
+44 (0)1223 274432 or 274431
- PatientLine
*801 (*from patient bedside telephones only*)
- E mail
pals@addenbrookes.nhs.uk
- Mail
PALS, Box No 53
Addenbrooke's Hospital
Hills Road, Cambridge, CB2 2QQ

Chaplaincy and Multi-Faith Community

- Telephone
+44 (0)1223 217769
- E mail
derek.fraser@addenbrookes.nhs.uk
- Mail
The Chaplaincy, Box No 105
Addenbrooke's Hospital
Hills Road, Cambridge, CB2 2QQ

MINICOM System ("type" system for the hard of hearing)

- Telephone
+44 (0)1223 274604

Access Office (travel, parking & security information)

- Telephone
+44 (0)1223 586969

What should I do with this form?

Thank you for taking the trouble to read this information sheet. If you wish to sign it and retain a copy for your own records, please do so below.

If you would like a copy of this form to be filed in your hospital records for future reference, please let your Urologist or Specialist Nurse know. If you do, however, decide to proceed with the scheduled procedure, you will be asked to sign a separate consent form which will be filed in your hospital notes and you will, in addition, be provided with a copy of the form if you wish.

I have read this information sheet and I accept the information it provides.

Signature.....Date.....

How can I get information in alternative formats?

Please ask if you require this information in other languages, large print or audio format: 01223 216032 or patient.information@addenbrookes.nhs.uk

- Polish** Informacje te można otrzymać w innych językach, w wersji dużym drukiem lub audio. Zamówienia prosimy składać pod numerem: 01223 216032 lub wysyłając e-mail: patient.information@addenbrookes.nhs.uk
- Portuguese** Se precisar desta informação num outro idioma, em impressão de letras grandes ou formato áudio por favor telefone para o 01223 216032 ou envie uma mensagem para: patient.information@addenbrookes.nhs.uk
- Russian** Если вам требуется эта информация на другом языке, крупным шрифтом или в аудиоформате, пожалуйста, обращайтесь по телефону 01223 216032 или на вебсайт patient.information@addenbrookes.nhs.uk
- Cantonese** 若你需要此信息的其他語言版本、大字體版或音頻格式，請致電 01223 216032 或發郵件到：patient.information@addenbrookes.nhs.uk
- Turkish** Bu bilgiyi diğer dillerde veya büyük baskılı ya da sesli formatta isterseniz lütfen su numaradan kontak kurun: 01223 216032 veya aşağıdaki adrese e-posta gönderin: patient.information@addenbrookes.nhs.uk

Bengali

এই তথ্য বাংলায়, বড় অক্ষরে বা অডিও টেপে পোতে চাইলে দয়া করে 01223 216032 নম্বরে ফোন করুন
বা patient.information@addenbrookes.nhs.uk ঠিকানায় ই-মেইল করুন।



Addenbrooke's is smoke-free. You cannot smoke anywhere on the site. Smoking increases the severity of some urological diseases and increases the risk of post-operative complications. For advice on quitting, contact your GP or the NHS smoking helpline free on 0800 169 0 169

Document history

Author(s)	Nikesh Thiruchelvam (on behalf of the Consultant Urologists)
Department	Department of Urology, Box No 43 Addenbrooke's Hospital Cambridge University Hospitals NHS Foundation Trust Hills Road Cambridge, CB2 2QQ www.addenbrookes.org.uk
Contact number	01223 216575
Fax number	01223 216069
Dept website	www.camurology.org.uk
First published	May 2005
Review date	April 2014
File name	Artificial urinary sphincter in men
Version number	6.0
Ref	116/Urol_04_11